

## 60 Market Center Dr. #103, Collierville, TN 38017

O: 901.316.5752 TP: 855.344.8724 F: 901.316.5760 TF: 844.588.5560 beneverepharmacy.com

| roday's date: |                 |
|---------------|-----------------|
| → NEW PATIENT | CURRENT PATIENT |

## **HEPATITIS B REFERRAL FORM**

|   | PATIENT INFORMATION   |   | PR   | ESCRIBER INFORMAT   | TION       |          |
|---|---|---|--|---|------------|----------|
| atient Name   |   |   | Prescriber Name  | DEA#  |            |          |
| ddress  |   |   | NPI#   | Tax ID_   |            |          |
| ty  | State   | Zip   | Medicaid ID #  |   |            |          |
| •   | Cell  | •   |  |   |            |          |
|   | SSN   |   |  | State   |            |          |
|   |   |   |  | Fax   |            |          |
| •   | Height:   |   |  | . <b>u</b> n  |            |          |
| ationic vivolgina.                                      |   |   | R MEDICAID INFORMA   |   |            |          |
|   |   |   |  |   |            |          |
| ,   |   |   | - ·  | olicable)   |            | roup     |
| surance Phone   |   |   |  |   |            |          |
| · · · · · · · · · · · · · · · · · · ·                   | e: Company  |   |  |   |            |          |
|   | RXBIN#  |   |  |   |            |          |
|   | COMPLETE OR FAX FRONT AN  | D BACK COPIES OF INSUR  | ANCE AND PRESCRIPTION BENEFIT  | CARDS IF AVAILABLE  |            |          |
|   |   | CLINICAL IN   | FORMATION  |   |            |          |
| agnosis: 🗆 B19.10 Chi                                   | ronic Hepatitis B without hepatic con   | na □ Other  |  | Diagn   | osis date: |          |
| •   | /labs   |   |  |   |            |          |
|   | men/Duration:)  |   |  |   |            |          |
| o-infection? 🗆 HCV 🗆                                    | HIV If applicable, please send all clinical infor   | mation pertinent to the patient   | 's co-infection. Scr D   | ate Pregnancy T   | est □ No □ | Yes □ N  |
|   | PLEA  | ASE ATTACH A LIST OF PATI   | ENT'S CURRENT MEDICATIONS.   |   |            |          |
| DRUG NAME   | DOSE  |   | DIDECTIONS FOR I   | ICE.  | QTY.       | REFILL   |
| DRUG NAME   |   |   | DIRECTIONS FOR U   | 9E  | uii.       | NEFILL   |
|   | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension  | ☐ Take (1) one tab<br>least 2 hours a   | plet daily on an empty stoma<br>fter a meal and 2 hours befo   | ch. Should be taken at  | UIT.       | NETTEE   |
|   | □ 0.5 mg tablet □ 1 mg tablet   | least 2 hours a   | olet daily on an empty stoma<br>fter a meal and 2 hours befo<br>hours after a meal and 2 ho  | ch. Should be taken at re the next meal.  Should be urs before the next meal.   |            | NEFILE   |
| □ Baraclude   | □ 0.5 mg tablet □ 1 mg tablet   | least 2 hours a   | olet daily on an empty stoma<br>fter a meal and 2 hours befo   | ch. Should be taken at re the next meal.  Should be urs before the next meal.   |            | NEFIEL   |
| □ Baraclude   | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet  | least 2 hours at taken at least 2  *** Dose must be adjust  Take 100 mg da  | olet daily on an empty stoma<br>fter a meal and 2 hours befo<br>hours after a meal and 2 ho<br>red for renal function. Please provide<br>ily with or without food.   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.   |            | NEFIEL   |
| □ Baraclude   | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension  | least 2 hours at taken at least 2 **** Dose must be adjust  | olet daily on an empty stoma<br>fter a meal and 2 hours befo<br>hours after a meal and 2 ho  | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.   |            | NETICE.  |
| □ Baraclude   | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution  | least 2 hours a   | olet daily on an empty stoma<br>fter a meal and 2 hours befo<br>hours after a meal and 2 ho<br>ted for renal function. Please provide<br>ily with or without food.   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.   |            | NETICE.  |
| □ Baraclude   | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet  | least 2 hours at taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da   | olet daily on an empty stoma fter a meal and 2 hours beforman hours after a meal and 2 hours after a meal function. Please provide the for renal function. Please provide the formal function. Please provide the formal function.   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.   |            | NETICE . |
| □ Baraclude   | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution  | least 2 hours at taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da   | olet daily on an empty stoma<br>fter a meal and 2 hours befo<br>hours after a meal and 2 ho<br>ted for renal function. Please provide<br>ily with or without food.   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.   |            | NETICE.  |
| □ Baraclude □ Epivir HBV □ Hepsera®                     | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution  | least 2 hours at taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da   | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide the for renal function. | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.   |            | NETICE . |
| Baraclude Epivir HBV Hepsera®                           | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution □ 10 mg tablet   | least 2 hours and taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da  *** Dose must be adjust  Take 1 tablet da   | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide the for renal function. | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.   |            | N.F.IC.  |
| □ Baraclude □ Epivir HBV □ Hepsera® □ Vemlidy®          | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution □ 10 mg tablet □ 25 mg tablet □ 300mg tablet □ 250mg tablet  | least 2 hours and taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da  *** Dose must be adjust  Take 1 tablet da  Take 1 tablet da  Take 300 mg da                                 | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide a filly with or without food and the form of the food and the food and the food and the food.   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.  current Scr and laboratory date.                                   |            | N.F.IL.  |
| □ Baraclude □ Epivir HBV □ Hepsera® □ Vemlidy®          | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution □ 10 mg tablet □ 25 mg tablet □ 300mg tablet □ 250mg tablet □ 200mg tablet   | least 2 hours and taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da  *** Dose must be adjust  Take 1 tablet da  Take 1 tablet da  Take 300 mg da                                 | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide ally with or without food and for renal function. Please provide ally with food   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.  current Scr and laboratory date.                                   |            |          |
|   | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution □ 10 mg tablet □ 25 mg tablet □ 300mg tablet □ 250mg tablet  | least 2 hours and taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da  *** Dose must be adjust  Take 1 tablet da  Take 1 tablet da  Take 300 mg da                                 | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide a filly with or without food and the form of the food and the food and the food and the food.   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.  current Scr and laboratory date.                                   |            |          |
| □ Baraclude □ Epivir HBV □ Hepsera® □ Vemlidy®          | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution □ 10 mg tablet □ 25 mg tablet □ 300mg tablet □ 250mg tablet □ 200mg tablet □ 150mg tablet                                      | least 2 hours and taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da  *** Dose must be adjust  Take 1 tablet da  Take 1 tablet da  Take 300 mg da                                 | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide ally with or without food and for renal function. Please provide ally with food   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.  current Scr and laboratory date.                                   |            |          |
| □ Baraclude □ Epivir HBV □ Hepsera® □ Vemlidy®          | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution □ 10 mg tablet □ 25 mg tablet □ 300mg tablet □ 250mg tablet □ 200mg tablet □ 150mg tablet                                      | least 2 hours and taken at least 2  *** Dose must be adjust  Take 100 mg da  *** Dose must be adjust  Take 1 tablet da  *** Dose must be adjust  Take 1 tablet da  *** Dose must be adjust  Take 300 mg da  *** Dose must be adjust | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide hilly with or without food and the for renal function. Please provide hilly with food hilly with or without food.  | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.  current Scr and laboratory date.                                   |            |          |
| □ Baraclude □ Epivir HBV □ Hepsera® □ Vemlidy® □ Viread | □ 0.5 mg tablet □ 1 mg tablet □ 0.05 mg/mL oral suspension □ 100 mg tablet □ 5 mg/mL oral solution □ 10 mg tablet □ 25 mg tablet □ 250mg tablet □ 250mg tablet □ 200mg tablet □ 150mg tablet □ 150mg tablet □ 40mg/gm oral powder | least 2 hours and taken at least 2 **** Dose must be adjust    Take 100 mg da    **** Dose must be adjust    Take 1 tablet da    **** Dose must be adjust    Take 1 tablet da    **** Dose must be adjust    MD's Office, remaining | plet daily on an empty stoma fter a meal and 2 hours befor hours after a meal and 2 hours befor hours after a meal and 2 hours after a meal function. Please provide ally with or without food and for renal function. Please provide ally with food ally with or without food.   | ch. Should be taken at re the next meal.  Should be urs before the next meal.  current Scr and laboratory date.  current Scr and laboratory date.  current Scr and laboratory date. |            |          |